


[Accedi / Registrati \(/user/login\)](/user/login)[Invia \(https://susy.mdpi.com/user/manuscripts/upload?journal=molecules\)](https://susy.mdpi.com/user/manuscripts/upload?journal=molecules)

### Cerca articoli :

### AvanzateRicerca

[Riviste \(/about/journals\)](/about/journals) / [Molecole \(/journal/molecules\)](/journal/molecules) / [Tomo 27 \(/1420-3049/27\)](/1420-3049/27) / [Problema 17 \(/1420-3049/27/17\)](/1420-3049/27/17) / [10.3390/molecules27175405](https://doi.org/10.3390/molecules27175405) /



*molecules*

[\(/journal/molecules\)](/journal/molecules)

Invia a questa rivista  
([https://susy.mdpi.com/user/manuscripts/upload?form%5Bjournal\\_id%5D%3D1](https://susy.mdpi.com/user/manuscripts/upload?form%5Bjournal_id%5D%3D1))

Recensione per questo Journal  
(<https://susy.mdpi.com/volunteer/journals/review>)

**Modifica un numero speciale**  
(</journalproposal/sendproposalspecialissue/molecules>)

► **Menù articolo**



**Maria José U. Ferreira**

[Iscriviti a SciFeed \(/1420-3049/27/17/5405/scifeed\\_display\)](#)

[Articoli consigliati](#)

[Collegamenti a informazioni correlate](#) v

[Altro per autori Collegamenti](#) v

**Visualizzazioni articolo** 121476

**Citazioni** 2




**Sommario** ^

- [Astratto](#)
- [Introduzione](#)
- [Risultati e discussione](#)
- [Materiali e metodi](#)
- [Conclusioni](#)
- [Materiali supplementari](#)
- [Contributi dell'autore](#)
- [Finanziamento](#)
- [Dichiarazione del comitato di revisione istituzionale](#)
- [Dichiarazione di consenso informato](#)
- [Dichiarazione sulla disponibilità dei dati](#)
- [Conflitto di interessi](#)
- [Riferimenti](#)

# Effetto degradativo di Nattokinase sulla proteina della punta di SARS-CoV-2

Condividere

di

 [\(/toggle desktop layout cookie\)](#)  

 **Takashi Tanikawa** (<https://sciprofiles.com/profile/1691959>)<sup>1,\*</sup>  (mailto:please\_login),

[Aiuto](#)

 **Yuka Kiba** (<https://sciprofiles.com/profile/author/YmdjSVRjSkpERWVQTHBkQ1dQUDI>)<sup>2,†</sup>,

[Citare](#)

 **Giacomo Yu** (<https://sciprofiles.com/profile/2378347>)<sup>3</sup>,

 **Kate Hsu** (<https://sciprofiles.com/profile/2378314>)<sup>3</sup>,

 **Shinder Chen** (<https://sciprofiles.com/profile/author/V0FuK1JjY3JKczVRUIZGK0EMOI>)<sup>3,†</sup>,

Discuti in

[Scienze](#)

[\(https://s](#)

[groups/p](#)



 **Ayako Ishii** (<https://sciprofiles.com/profile/2404433>)<sup>4</sup>,

 **Takami Yokogawa** (<https://sciprofiles.com/profile/2404391>)<sup>2</sup>  (<https://orcid.org/0000-0001-9713-8505>),



[Approvare](#)

 **Ryuichiro Suzuki** (<https://sciprofiles.com/profile/332434>)<sup>5</sup>,

 **Yutaka Inoue** (<https://sciprofiles.com/profile/531611>)<sup>1</sup>  (<https://orcid.org/0000-0003-3419-343X>) e

[Commento](#)

 **Masashi Kitamura** (<https://sciprofiles.com/profile/1186307>)<sup>2,\*</sup>  (mailto:please\_login)

<sup>1</sup> Laboratorio di Gestione della Nutri-Farmacoterapia, Scuola di Farmacia, Facoltà di Farmacia e Scienze Farmaceutiche, Università di Josai, Saitama 350-0295, Giappone

<sup>2</sup> Laboratorio di Farmacognozia, Scuola di Farmacia, Facoltà di Farmacia e Scienze Farmaceutiche, Università di Josai, Saitama 350-0295, Giappone

<sup>3</sup> Contek Life Science Co., Ltd., città di Taipei 100007, Taiwan

<sup>4</sup> CellMark Giappone, Tokyo 102-0071, Giappone

<sup>5</sup> Laboratorio di Prodotti Naturali e Fitochimica, Dipartimento di Scienze Farmaceutiche, Facoltà di Farmacia e Scienze Farmaceutiche, Josai University, Saitama 350-0295, Giappone

\* Autori a cui indirizzare la corrispondenza.

† Questi autori hanno contribuito ugualmente a questo lavoro.

*Molecole* **2022**, 27 (17), 5405; <https://doi.org/10.3390/molecules27175405>  
(<https://doi.org/10.3390/molecules27175405>)

**Ricevuto: 14 luglio 2022 / Revisionato: 17 agosto 2022 / Accettato: 23 agosto 2022 /  
Pubblicato: 24 agosto 2022**

(Questo articolo appartiene al numero speciale **Composti bioattivi negli alimenti fermentati sani** ( /journal/molecules/special\_issues/healthy\_fermented\_food ) )

res.com/molecules/molecules-27-05405/article\_deploy/html/images/molecules-27-05405-g001a.png?1662454982) (https://pub.mdpi-res.com/molecules/molecules-27-05405/article\_deploy/html/images/molecules-27-05405-g001b.png?1662454978) (https://pub.mdpi-res.com/molecules/molecules-27-05405/article\_deploy/html/images/molecules-27-05405-g002.png?1662454986)

Esamina i rapporti (/1420-3049/27/17/5405/review\_report)

Versioni Note (/1420-3049/27/17/5405/notes)

## Astratto

La malattia da coronavirus 2019 (COVID-19), causata dalla sindrome respiratoria acuta grave coronavirus 2 (SARS-CoV-2), è emersa come una pandemia e ha inflitto enormi danni alla vita delle persone e all'economia di molti paesi in tutto il mondo. Tuttavia, gli agenti terapeutici contro SARS-CoV-2 rimangono poco chiari. SARS-CoV-2 ha una proteina spike (proteina S) e la scissione della proteina S è essenziale per l'ingresso virale. Nattokinase è prodotto da *Bacillus subtilis* var. *nattoed* è benefico per la salute umana. In questo studio, abbiamo esaminato l'effetto della nattokinase sulla proteina S di SARS-CoV-2. Quando i lisati cellulari trasfettati con la proteina S sono stati incubati con nattokinase, la proteina S è stata degradata in modo dipendente dalla dose e dal tempo. L'analisi dell'immunofluorescenza ha mostrato che la proteina S sulla superficie cellulare è stata degradata quando la nattokinasi è stata aggiunta al terreno di coltura. Pertanto, i nostri risultati suggeriscono che la nattokinase mostra il potenziale per l'inibizione dell'infezione da SARS-CoV-2 tramite la degradazione della proteina S.

**Parole chiave:** SARS-CoV-2 (/search?q=SARS-CoV-2) ; nattokinasi (/search?q=nattokinase) ; COVID 19 (/search?q=COVID-19)

## 1. Introduzione

Coronavirus disease 2019 (COVID-19), caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), has been spreading worldwide. The COVID-19 pandemic has affected over 437 million people and caused more than 6.3 million deaths (https://covid19.who.int/ (https://covid19.who.int/), accessed on 4 July 2022). The entry of SARS-CoV-2 into host cells is mediated by the transmembrane spike protein (S protein),

MDPI (1)

which forms homotrimers that extend from the viral envelope. The S protein is processed and activated by cellular proteases including transmembrane serine protein 2 (TMPRSS2), cathepsin, and furin. It comprises two functional subunits, S1 and S2; the S1 subunit of SARS-CoV-2 initiates virus-receptor binding by interacting with the human host cell receptor angiotensin-converting enzyme 2 (ACE2), and the S2 subunit participates in viral fusion with the target cell, allowing viral entry [1]. The receptor-binding domain (RBD) in the S1 subunit is responsible for binding to ACE2. S protein cleavage occurs at the boundary between the S1 and S2 subunits.

Currently, many countries are involved in the development of vaccines to protect against SARS-CoV-2 infection, thus, the number of SARS-CoV-2 infections has decreased. However, numerous variants of SARS-CoV-2, including strains with mutated vaccine target epitopes, have been reported [2,3]. Vaccination may not completely protect against SARS-CoV-2 infection because the number of patients with COVID-19 is increasing after vaccination. Therefore, it is important to develop novel treatments for SARS-CoV-2 infections.

Natto is a popular traditional Japanese food made from soybeans fermented by *Bacillus subtilis* var. *natto*. Nattokinase is found in natto [4] and is one of the most important extracellular enzymes produced by *B. subtilis* var. *natto* [5]. Nattokinase consists of 275 amino acids and is approximately 28 kDa [6,7]. Nattokinase inactivates plasminogen activator inhibitor-1 and increases fibrinolysis [8]. It also decreases the plasma levels of fibrinogen, factor VII, cytokines, and factor VIII [9]. Nattokinase has the highest clot-dissolving potency among naturally known anticoagulants [10]. A clinical trial demonstrated that oral consumption of nattokinase was not associated with any adverse effects [11]. Thus, nattokinase is now considered an efficient, secure, and economical enzyme that has drawn central attention in thrombolytic drug studies [12,13]. In addition, nattokinase is used in the treatment of some tumors [14,15].

A recent study revealed that natto extract inhibits bovine herpesvirus 1 (BHV-1) and SARS-CoV-2 infection [16]. These results indicate that natto extract protease might be effective against SARS-CoV-2 infection. In this study, we aimed to investigate whether the inhibition of SARS-CoV-2 infection by natto extract is caused by nattokinase derived from *B. subtilis* var. *natto*.


## 2. Results and Discussion

### 2.1. Degradative Effects of Nattokinase on Spike Protein of SARS-CoV-2 In Vitro


We first investigated whether nattokinase in natto extract could degrade SARS-CoV-2 S protein. The S protein of SARS-CoV-2 plays an important role in the ACE2 receptor of the host cell during the early stages of infection [17]. After mixing the S protein expression cell lysate with a 4-fold dilution series of nattokinase (32 µg/mL, 8 µg/mL, 2 µg/mL, 500 ng/mL, 125 ng/mL, 31.25 ng/mL, and 7.8125 ng/mL), Western blotting was performed. The full length of S protein (S1 and S2 subunits) and S2 subunit appeared as bands when S protein

MDPI (l)

expression cell lysate was incubated with D-PBS at nattokinase concentrations of 500 ng/mL, 125 ng/mL, 31.25 ng/mL, and 7.8125 ng/mL (**Figure 1A**). Next, we examined whether nattokinase degrades the S protein in a time-dependent manner. The lysate was then incubated with 1 µg/mL nattokinase for 10–180 min. The S protein of SARS-CoV-2 was degraded by nattokinase after 60–180 min of incubation, but not after 10 and 30 min of incubation (**Figure 1B**). Thus, nattokinase degraded S protein in a dose- and time-dependent manner.




**Figure 1.** (A) Degradative effects of nattokinase in dose-dependent manner. Serially diluted nattokinase (32 µg/mL, 8 µg/mL, 2 µg/mL, 500 ng/mL, 125 ng/mL, 31.25 ng/mL, and 7.8125 ng/mL) were mixed with S protein expression cell lysate and incubated. Full length of S protein (S1 and S2 subunits) and S2 subunit were detected as upper and lower bands, respectively. Ratio of total S was indicated as the relative quantity of S protein (S protein + S2 protein). (B) Degradative effects of nattokinase in time-dependent manner. S protein expression cell lysate was incubated with 1 µg/mL nattokinase for 0, 10, 30, 60, 120, and 180 min. (C) Effects of heating treatment or protease inhibitors. Lane 1: HEK293 lysate; lane 2: HEK293 lysate (S protein); lane 3: HEK293 (S protein) + nattokinase (5 µg/mL); lane 4: HEK293 (S protein) + nattokinase (5 µg/mL) + Protease inhibitor I; lane 5: HEK293 (S protein) + nattokinase (5 µg/mL) + Protease inhibitor III; lane 6: HEK293 (S protein) + heat-treated nattokinase (5 µg/mL). (D) Degradative effect on RBD of S protein and ACE2. RBD of S protein and ACE2 coding plasmids were transfected with HEK293 cells, respectively. Cell lysates were incubated with nattokinase (7.5 µg/mL) and heat-treated nattokinase (7.5 µg/mL) and Western blotting was performed.

 To confirm whether the degradative effect of nattokinase is due to enzymatic activity, nattokinase was treated with heating or a protease inhibitor cocktail. When nattokinase was heated at 100 °C for 5 min, the degradative effect of nattokinase was lost (**Figure 1C**, lane 6). Furthermore, the loss of the S protein bands by nattokinase was blocked when protease inhibitors were added (**Figure 1C**, lanes 4 and 5). Compared with protein inhibitor cocktail I, protein cocktail III, which consisted of AEBSF HCl (4-(2-Aminoethyl) benzenesulfonyl fluoride hydrochloride), aprotinin, which is an irreversible serine protease inhibitor, and leupeptin, which is a cysteine-protease, clearly blocked nattokinase activity. Nattokinase has the same conserved amino acids, Ser-His-Asp (Asp<sup>32</sup>, His<sup>64</sup>, and Ser<sup>221</sup>), which are members of the subtilisin family of serine proteases [6,18]. The crystal structure of nattokinase is nearly identical to that of subtilisin E from *B. subtilis* DB104 [19]. This result is consistent with that of a previous report that nattokinase is a serine protease. We also assessed the degradative effects of nattokinase using cell lysates expressing the RBD and ACE2. When 7.5 µg/mL of nattokinase and cell lysate were incubated, the bands of RBD and ACE2 were lost (**Figure 1D**).

## 2.2. Degradative Effects of Nattokinase on Spike Protein of SARS-CoV-2 on the Transfected Cell Surface

Next, we examined whether nattokinase degrades the S protein on the transfected cell surface. The S protein was transfected with the HEK293 cells. The transfected cells were incubated with nattokinase for 9 h. The S protein on the cell surface was detected using an anti-S protein antibody without cell permeabilization (**Figure 2A**). The S protein was detected in the transfected cells. When transfected cells were treated with nattokinase, the S protein on the cell surface decreased. When cells were treated with 25 µg/mL and 2.5 µg/mL nattokinase, the ratio of S protein-positive area to nucleus-positive area decreased by approximately 0.3 and 0.7, respectively (**Figure 2B**). The degradative effect of nattokinase was observed when there was no cytotoxicity (**Figure 2C**). Western blotting analysis showed that the quantity of total S protein did not change among nattokinase and control treatments (**Supplemental Figure; Figure S1**). These results indicate that nattokinase would degrade the S protein of SARS-CoV-2 in the non-toxic concentration range.



**Figure 2.** (A) Degradative effect of nattokinase on S protein on the cell surface. Spike-pcDNA3.1 was transfected with HEK2993 cells and incubated for 9 h. After incubation, nattokinase (25 and 2.5 µg/mL) were added to culture medium and further incubated for 13 h. Cells were fixed and immunofluorescent analysis was performed. S protein on the cell surface was stained with anti-spike protein antibody (Red) and nucleus was stained with DAPI (Blue). (B) Ratio of S protein area to nucleus positive area. Three images per sample were captured and S protein/nucleus positive areas were calculated. Data are shown as mean + SD, and  $p$ -value was determined by one-way analysis of variance (ANOVA) with Tukey's post-hoc test using R software (R-3.3.3 for windows) (\*\*  $p < 0.01$ ; \*\*\*  $p < 0.001$ ). (C) Cell viability was evaluated by MTT assay. Indicated nattokinase was added to culture medium and incubated for 13 h; MTT assay was performed.

In this study, we showed that the protease activity of nattokinase contributes to the degradation of S protein. Nattokinase has a degrading effect on not only S proteins but also ACE2 in host cells. The protease specificity of nattokinase would be low, because GAPDH, a housekeeping protein, was also degraded simultaneously in the in-vitro evaluation of nattokinase mixed with cell lysate (**Supplemental Figure; Figure S2**). On the other hand, when added to cells, it does not show any effect on cell viability and is expected to act as a protective agent on the cell surface. Further analysis of the degradation products of nattokinase using mass spectrometry is needed for understanding the proteolysis effects.

Nattokinase possesses the potent degradation activity for SARS-CoV-2 S protein and has also been shown to exert anti-atherosclerotic, lipid-lowering, antihypertensive, antithrombotic, fibrinolytic, neuroprotective, antiplatelet, and anticoagulant effects [20]. Patients with hypertension and cardiovascular comorbidities can easily get very sick from COVID-19 [21]. Due to the emergence of numerous variants of SARS-CoV-2 including strains with mutated vaccine target epitopes, vaccination alone may not completely protect against



SARS-CoV-2 infection. Nattokinase and natto extracts have the potential to be developed as a new generation of drug for the prevention and treatment of COVID-19.

 [\(toggle desktop layout cookie\)](#)  

### 3. Materials and Methods

#### 3.1. Materials

Nattokinase was obtained from Contek Life Science Co., Ltd. (Taipei City, Taiwan). The nattokinase activity was 60,000 FU/g (FU, fibrinolysis unit). Protease inhibitor cocktail sets I and III were purchased from FUJIFILM Wako Pure Chemical Corporation (Osaka, Japan). The expression plasmid (pcDNA3.1-SARS2-Spike C9 with tag at the C-terminal, pcDNA3.1-hACE2, and pcDNA3-SARS-CoV-2-S-RBD-sfGFP) was purchased from Addgene (Watertown, MA, USA). HEK293 (JCRB9068) cells were obtained from the JCRB Cell Bank (Osaka, Japan).

#### 3.2. Cell Culture and Western Blotting

HEK293 cells were cultured at a density of  $3.5 \times 10^5$  cells/mL in DMEM supplemented with 10% FBS, L-glutamine, 100 U/mL penicillin, and 100  $\mu$ g/mL streptomycin overnight. The cells were transfected with each plasmid (pcDNA3.1-SARS2-Spike, pcDNA3-SARS-CoV-2-S-RBD-sfGFP, or pcDNA3.1-hACE2) and incubated for 22 h. After incubation, the cultured cells were scraped and washed with ice-cold Dulbecco's phosphate-buffered saline (D-PBS). Cell counting was performed and xTractor buffer (Takara Bio Inc., Shiga, Japan) was added to the cell precipitate. The cell lysates were centrifuged at  $1300 \times g$  for 10 min at 4 °C and the supernatant was transferred to new tubes and stored at -80 °C until use. The protein concentration was determined by bicinchoninic acid (BCA) protein assay using a BCA assay kit (Takara). Ten microliters of nattokinase and 10  $\mu$ L of cell lysate (1  $\mu$ g/mL) were incubated at 37 °C for 1 h. When the effects of protease inhibitors were used, protease inhibitor cocktail sets I and III were diluted 10-fold with D-PBS, and a 10  $\mu$ L protease inhibitor cocktail solution was added to the mixture of nattokinase and cell lysate. Equal volumes of the reaction mixture were loaded and Western blotting was performed. The primary antibodies included anti-rhodopsin (C9) mouse monoclonal antibody (1D4) (Santa Cruz Biotechnology, Dallas, TX, USA), anti-GAPDH mouse monoclonal antibody (FUJIFILM Wako), anti-GFP tag mouse monoclonal antibody (Proteintech, Rosemont, IL, USA), and anti-ACE2 antibody (Proteintech). Secondary antibodies include HRP-conjugated goat anti-mouse antibody (Proteintech).

#### 3.3. Immunofluorescence Assay

HEK293 cells were cultured at a density of  $3.5 \times 10^5$  cells/mL in an 8-well chamber in DMEM supplemented with 10% FBS, L-glutamine, 100 U/mL penicillin, and 100  $\mu$ g/mL streptomycin. The cells were transfected with pcDNA3.1-SARS2-Spike and incubated for 9 h. After incubation, the cells were treated with the samples, incubated for 13 h, and fixed in 4% paraformaldehyde for 30 min. After incubation with SARS-CoV/SARS-CoV-2 spike

monoclonal antibody (1A9) (GeneTex, CA, USA) for 1 h, it was incubated with Cy3-conjugated goat anti-mouse antibody for 1 h. The slides were stained with DAPI Fluoromount-G and observed using a fluorescence microscope (BZ-X710, Keyence, Osaka, Japan). S protein-positive and nucleus-positive areas were calculated using BZ-X710 attached analysis software (BZ-X Analyzer). Cell viability was assessed using the 3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. The cells were cultivated in 24-well culture plates. After incubation at 37 °C for 24 h, samples were added to each well and incubated for another 13 h. Cells were suspended in 500 µL of DMEM containing 500 µg/mL MTT. After incubation for 3 h at 37 °C, 500 µL isopropanol containing 4 mM HCl was added to dissolve MTT formazan. The absorbance was measured at 570 nm using a microplate reader.

#### 4. Conclusions

In this study, we demonstrated that nattokinase, a serine protease, degrades the S protein of SARS-CoV-2. To investigate whether nattokinase contained in natto extract could inhibit SARS-CoV-2 infection, we analyzed S protein degradation by mixing the S protein expression cell lysate and nattokinase in a dose- and time-dependent manner. The RBD of the S protein binds to the membrane-distal portion of the ACE2 protein. Natto extract has been reported to inhibit SARS-CoV-2 infection in Vero E6 cells via RBD degradation [16]. We demonstrated that S protein degradation by nattokinase was blocked by heat or protein-inhibitor treatments. Our data suggest that the protease activity of nattokinase plays a crucial role in S protein degradation. Taken together, these findings support the notion that the inhibition of SARS-CoV-2 infection by natto extract was due to S protein degradation by nattokinase. Thus, our data indicated that nattokinase and natto extracts have potential effects on the inhibition of SARS-CoV-2 host cell entry via S protein degradation.

#### Supplementary Materials

The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/molecules27175405/s1> (<https://www.mdpi.com/article/10.3390/molecules27175405/s1>), Figure S1: Effect of nattokinase addition to S protein expressing cell culture medium was evaluated by Western blotting; Figure S2: Degradative effects of GAPDH in vitro.

#### Author Contributions

Conceptualization and supervision, T.T., J.Y., K.H., S.C., A.I., T.Y., R.S., Y.I. and M.K.; Methodology, all authors; Western blot and immunofluorescent analyses, T.T., Y.K. and M.K.; writing—original draft preparation, T.T. and M.K.; writing—review and editing, all authors. All authors have read and agreed to the published version of the manuscript.

## Institutional Review Board Statement

Not applicable.

## Informed Consent Statement

Not applicable.

## Data Availability Statement

The data used to support these findings have been included in this article. Additional information is available from the corresponding authors upon request.

## Conflicts of Interest

The authors declare no conflict of interest.

## References

1. Alsobaie, S. Understanding the Molecular Biology of SARS-CoV-2 and the COVID-19 Pandemic: A Review. *Infect. Drug Resist.* **2021**, *14*, 2259–2268. [Google Scholar ([https://scholar.google.com/scholar\\_lookup?title=Understanding+the+Molecular+Biology+of+SARS-CoV-2+and+the+COVID-19+Pandemic:+A+Review&author=Alsobaie,+S.&publication\\_year=2021&journal=Infect.+Drug+Resist.&volume=14&pages=2259%E2%80%932268&doi=10.2147/IDR.S306441&pmid=34163190](https://scholar.google.com/scholar_lookup?title=Understanding+the+Molecular+Biology+of+SARS-CoV-2+and+the+COVID-19+Pandemic:+A+Review&author=Alsobaie,+S.&publication_year=2021&journal=Infect.+Drug+Resist.&volume=14&pages=2259%E2%80%932268&doi=10.2147/IDR.S306441&pmid=34163190))] [CrossRef (<https://doi.org/10.2147/IDR.S306441>)] [PubMed (<http://www.ncbi.nlm.nih.gov/pubmed/34163190>)]
2. Bayati, A.; Kumar, R.; Francis, V.; McPherson, P.S. SARS-CoV-2 infects cells after viral entry via clathrin-mediated endocytosis. *J. Biol. Chem.* **2021**, *296*, 100306. [Google Scholar ([https://scholar.google.com/scholar\\_lookup?title=SARS-CoV-2+infects+cells+after+viral+entry+via+clathrin-mediated+endocytosis&author=Bayati,+A.&author=Kumar,+R.&author=Francis,+V.&author=McPherson,+P.S.&publication\\_year=2021&journal=J.+Biol.+Chem.&volume=296&pages=100306&doi=10.1016/j.jbc.2021.100306&pmid=33476648](https://scholar.google.com/scholar_lookup?title=SARS-CoV-2+infects+cells+after+viral+entry+via+clathrin-mediated+endocytosis&author=Bayati,+A.&author=Kumar,+R.&author=Francis,+V.&author=McPherson,+P.S.&publication_year=2021&journal=J.+Biol.+Chem.&volume=296&pages=100306&doi=10.1016/j.jbc.2021.100306&pmid=33476648))] [CrossRef (<https://doi.org/10.1016/j.jbc.2021.100306>)] [PubMed (<http://www.ncbi.nlm.nih.gov/pubmed/33476648>)]

Bestle, D.; Heindl, M.R.; Limburg, H.; Van Lam van, T.; Pilgram, O.; Moulton, H.; Stein, D.A.; Harges, K.; Eickmann, M.; Dolnik, O.; et al. TMPRSS2 and furin are both essential for proteolytic activation of SARS-CoV-2 in human airway cells. *Life Sci. Alliance* **2020**, *3*, e202000786. [Google Scholar

([https://scholar.google.com/scholar\\_lookup?](https://scholar.google.com/scholar_lookup?title=TMPRSS2+and+furin+are+both+essential+for+proteolytic+activation+of+SARS-CoV-2+in+human+airway+cells&author=Bestle,+D.&author=Heindl,+M.R.&author=Limburg,+H.&author=Van+Lam+van,+T.&author=Pilgram,+O.&author=Moulton,+H.&author=Stein,+D.A.&author=Harges,+K.&author=Eickmann,+M.&author=Dolnik,+O.&publication_year=2020&journal=Life+Sci.+Alliance&volume=3&pages=e202000786&doi=10.26508/lsa.202000786&pmid=32703818)

[title=TMPRSS2+and+furin+are+both+essential+for+proteolytic+activation+of+SARS-CoV-](https://scholar.google.com/scholar_lookup?title=TMPRSS2+and+furin+are+both+essential+for+proteolytic+activation+of+SARS-CoV-2+in+human+airway+cells&author=Bestle,+D.&author=Heindl,+M.R.&author=Limburg,+H.&author=Van+Lam+van,+T.&author=Pilgram,+O.&author=Moulton,+H.&author=Stein,+D.A.&author=Harges,+K.&author=Eickmann,+M.&author=Dolnik,+O.&publication_year=2020&journal=Life+Sci.+Alliance&volume=3&pages=e202000786&doi=10.26508/lsa.202000786&pmid=32703818)

[2+in+human+airway+cells&author=Bestle,+D.&author=Heindl,+M.R.&author=Limburg,+H.&author=Van+Lam+van,+T.&author=Pilgram,+O.&author=Moulton,+H.&author=Stein,+D.A.&author=Harges,+K.&author=Eickmann,+M.&author=Dolnik,+O.&publication\\_year=2020&journal=Life+Sci.+Alliance&volume=3&pages=e202000786&doi=10.26508/lsa.202000786&pmid=32703818">2+in+human+airway+cells&author=Bestle,+D.&author=Heindl,+M.R.&author=Limburg,+H.&author=Van+Lam+van,+T.&author=Pilgram,+O.&author=Moulton,+H.&author=Stein,+D.A.&author=Harges,+K.&author=Eickmann,+M.&author=Dolnik,+O.&publication\\_year=2020&journal=Life+Sci.+Alliance&volume=3&pages=e202000786&doi=10.26508/lsa.202000786&pmid=32703818](https://scholar.google.com/scholar_lookup?title=TMPRSS2+and+furin+are+both+essential+for+proteolytic+activation+of+SARS-CoV-2+in+human+airway+cells&author=Bestle,+D.&author=Heindl,+M.R.&author=Limburg,+H.&author=Van+Lam+van,+T.&author=Pilgram,+O.&author=Moulton,+H.&author=Stein,+D.A.&author=Harges,+K.&author=Eickmann,+M.&author=Dolnik,+O.&publication_year=2020&journal=Life+Sci.+Alliance&volume=3&pages=e202000786&doi=10.26508/lsa.202000786&pmid=32703818)] [CrossRef

(<https://doi.org/10.26508/lsa.202000786>)]

[CrossRef

[PubMed

(<http://www.ncbi.nlm.nih.gov/pubmed/32703818>)]

4. Sumi, H.; Hamada, H.; Tsushima, H.; Mihara, H.; Muraki, H. A novel fibrinolytic enzyme (nattokinase) in the vegetable cheese Natto; a typical and popular soybean food in the Japanese diet. *Experientia* **1987**, *43*, 1110–1111. [Google Scholar ([https://scholar.google.com/scholar\\_lookup?title=A+novel+fibrinolytic+enzyme+\(nattokinase\)+in+the+vegetable+cheese+Natto;+a+typical+and+popular+soybean+food+in+the+Japanese+diet&author=Sumi,+H.&author=Hamada,+H.&author=Tsushima,+H.&author=Mihara,+H.&author=Muraki,+H.&publication\\_year=1987&journal=Experientia&volume=43&pages=1110%E2%80%931111&doi=10.1007/BF01956052&pmid=3478223](https://scholar.google.com/scholar_lookup?title=A+novel+fibrinolytic+enzyme+(nattokinase)+in+the+vegetable+cheese+Natto;+a+typical+and+popular+soybean+food+in+the+Japanese+diet&author=Sumi,+H.&author=Hamada,+H.&author=Tsushima,+H.&author=Mihara,+H.&author=Muraki,+H.&publication_year=1987&journal=Experientia&volume=43&pages=1110%E2%80%931111&doi=10.1007/BF01956052&pmid=3478223))] [CrossRef (<https://doi.org/10.1007/BF01956052>)] [PubMed (<http://www.ncbi.nlm.nih.gov/pubmed/3478223>)]
5. Dabbagh, F.; Negahdaripour, M.; Berenjian, A.; Behfar, A.; Mohammadi, F.; Zamani, M.; Irajie, C.; Ghasemi, Y. Nattokinase: Production and application. *Appl. Microbiol. Biotechnol.* **2014**, *98*, 9199–9206. [Google Scholar ([https://scholar.google.com/scholar\\_lookup?title=Nattokinase:+Production+and+application&author=Dabbagh,+F.&author=Negahdaripour,+M.&author=Berenjian,+A.&author=Behfar,+A.&author=Mohammadi,+F.&author=Zamani,+M.&author=Irajie,+C.&author=Ghasemi,+Y.&publication\\_year=2014&journal=Appl.+Microbiol.+Biotechnol.&volume=98&pages=9199%E2%80%939206&doi=10.1007/s00253-014-6135-3&pmid=25348469](https://scholar.google.com/scholar_lookup?title=Nattokinase:+Production+and+application&author=Dabbagh,+F.&author=Negahdaripour,+M.&author=Berenjian,+A.&author=Behfar,+A.&author=Mohammadi,+F.&author=Zamani,+M.&author=Irajie,+C.&author=Ghasemi,+Y.&publication_year=2014&journal=Appl.+Microbiol.+Biotechnol.&volume=98&pages=9199%E2%80%939206&doi=10.1007/s00253-014-6135-3&pmid=25348469))] [CrossRef (<https://doi.org/10.1007/s00253-014-6135-3>)] [PubMed (<http://www.ncbi.nlm.nih.gov/pubmed/25348469>)]

6. Nakamura, T.; Yamagata, Y.; Ichishima, E. Nucleotide sequence of the Subtilisin NAT gene, aprN, of *Bacillus subtilis* (natto). *Biosci. Biotech. Bioch.* **1992**, *56*, 1869–1871. [Google Scholar ([https://scholar.google.com/scholar\\_lookup?title=Nucleotide+sequence+of+the+Subtilisin+NAT+gene,+aprN,+of+Bacillus+subtilis+\(natto\)&author=Nakamura,+T.&author=Yamagata,+Y.&author=Ichishima,+E.&publication\\_year=1992&journal=Biosci.+Biotech.+Bioch.&volume=56&pages=1869%E2%80%931871&doi=10.1271/bbb.56.1869&pmid=1369081](https://scholar.google.com/scholar_lookup?title=Nucleotide+sequence+of+the+Subtilisin+NAT+gene,+aprN,+of+Bacillus+subtilis+(natto)&author=Nakamura,+T.&author=Yamagata,+Y.&author=Ichishima,+E.&publication_year=1992&journal=Biosci.+Biotech.+Bioch.&volume=56&pages=1869%E2%80%931871&doi=10.1271/bbb.56.1869&pmid=1369081))] [CrossRef (<https://doi.org/10.1271/bbb.56.1869>)] [PubMed (<http://www.ncbi.nlm.nih.gov/pubmed/1369081>)]
7. Urano, T.; Ihara, H.; Umemura, K.; Suzuki, Y.; Oike, M.; Akita, S.; Tsukamoto, Y.; Suzuki, I.; Takada, A. The profibrinolytic enzyme subtilisin NAT purified from *Bacillus subtilis* Cleaves and inactivates plasminogen activator inhibitor type 1. *J. Biol. Chem.* **2001**, *276*, 24690–24696. [Google Scholar ([https://scholar.google.com/scholar\\_lookup?title=The+profibrinolytic+enzyme+subtilisin+NAT+purified+from+Bacillus+subtilis+Cleaves+and+inactivates+plasminogen+activator+inhibitor+type+1&author=Urano,+T.&author=Ihara,+H.&author=Umemura,+K.&author=Suzuki,+Y.&author=Oike,+M.&author=Akita,+S.&author=Tsukamoto,+Y.&author=Suzuki,+I.&author=Takada,+A.&publication\\_year=2001&journal=J.+Biol.+Chem.&volume=276&pages=24690%E2%80%9324696&doi=10.1074/jbc.M101751200](https://scholar.google.com/scholar_lookup?title=The+profibrinolytic+enzyme+subtilisin+NAT+purified+from+Bacillus+subtilis+Cleaves+and+inactivates+plasminogen+activator+inhibitor+type+1&author=Urano,+T.&author=Ihara,+H.&author=Umemura,+K.&author=Suzuki,+Y.&author=Oike,+M.&author=Akita,+S.&author=Tsukamoto,+Y.&author=Suzuki,+I.&author=Takada,+A.&publication_year=2001&journal=J.+Biol.+Chem.&volume=276&pages=24690%E2%80%9324696&doi=10.1074/jbc.M101751200))] [CrossRef (<https://doi.org/10.1074/jbc.M101751200>)]
8. Suzuki, Y.; Kondo, K.; Matsumoto, Y.; Zhao, B.Q.; Otsuguro, K.; Maeda, T.; Tsukamoto, Y.; Urano, T.; Umemura, K. Dietary supplementation of fermented soybean, natto, suppresses intimal thickening and modulates the lysis of mural thrombi after endothelial injury in rat femoral artery. *Life Sci.* **2003**, *73*, 1289–1298. [Google Scholar ([https://scholar.google.com/scholar\\_lookup?title=Dietary+supplementation+of+fermented+soybean,+natto,+suppresses+intimal+thickening+and+modulates+the+lysis+of+mural+thrombi+after+endothelial+injury+in+rat+femoral+artery&author=Suzuki,+Y.&author=Kondo,+K.&author=Matsumoto,+Y.&author=Zhao,+B.Q.&author=Otsuguro,+K.&author=Maeda,+T.&author=Tsukamoto,+Y.&author=Urano,+T.&author=Umemura,+K.&publication\\_year=2003&journal=Life+Sci.&volume=73&pages=1289%E2%80%931298&doi=10.1016/S0024-3205\(03\)00426-0](https://scholar.google.com/scholar_lookup?title=Dietary+supplementation+of+fermented+soybean,+natto,+suppresses+intimal+thickening+and+modulates+the+lysis+of+mural+thrombi+after+endothelial+injury+in+rat+femoral+artery&author=Suzuki,+Y.&author=Kondo,+K.&author=Matsumoto,+Y.&author=Zhao,+B.Q.&author=Otsuguro,+K.&author=Maeda,+T.&author=Tsukamoto,+Y.&author=Urano,+T.&author=Umemura,+K.&publication_year=2003&journal=Life+Sci.&volume=73&pages=1289%E2%80%931298&doi=10.1016/S0024-3205(03)00426-0))] [CrossRef ([https://doi.org/10.1016/S0024-3205\(03\)00426-0](https://doi.org/10.1016/S0024-3205(03)00426-0))]

Nattokinase decreases plasma levels of fibrinogen, factor VII, and factor VIII in human subjects. *Nutr. Res.* **2009**, *29*, 190–196. [Google Scholar

([https://scholar.google.com/scholar\\_lookup?](https://scholar.google.com/scholar_lookup?)

**title=Nattokinase+decreases+plasma+levels+of+fibrinogen,+factor+VII,+and+factor+VIII+in+human+subjects&author=Hsia,+C.H.&author=Shen,+M.C.&author=Lin,+J.S.&author=Wen,+Y.K.&author=Hwang,+K.L.&author=Cham,+T.M.&author=Yang,+N.C.&publication\_year=2009&journal=Nutr.+Res.&volume=29&pages=190%E2%80%93196&doi=10.1016/j.nutres.2009.01.009]** [CrossRef  
(<https://doi.org/10.1016/j.nutres.2009.01.009>)]

10. Weng, Y.; Yao, J.; Sparks, S.; Wang, K.Y. Nattokinase: An Oral Antithrombotic Agent for the Prevention of Cardiovascular Disease. *Int. J. Mol. Sci.* **2017**, *18*, 523. [Google Scholar

([https://scholar.google.com/scholar\\_lookup?](https://scholar.google.com/scholar_lookup?)  
**title=Nattokinase:+An+Oral+Antithrombotic+Agent+for+the+Prevention+of+Cardiovascular+Disease&author=Weng,+Y.&author=Yao,+J.&author=Sparks,+S.&author=Wang,+K.Y.&publication\_year=2017&journal=Int.+J.+Mol.+Sci.&volume=18&pages=523&doi=10.3390/ijms18030523]** [CrossRef  
(<https://doi.org/10.3390/ijms18030523>)] [Green Version (<https://www.mdpi.com/1422-0067/18/3/523/pdf>)]

11. Kim, J.Y.; Gum, S.N.; Paik, J.K.; Lim, H.H.; Kim, K.C.; Ogasawara, K.; Inoue, K.; Park, S.; Jang, Y.; Lee, J.H. Effects of nattokinase on blood pressure: A randomized, controlled trial. *Hypertens. Res.* **2008**, *31*, 1583–1588. [Google Scholar

([https://scholar.google.com/scholar\\_lookup?](https://scholar.google.com/scholar_lookup?)  
**title=Effects+of+nattokinase+on+blood+pressure:+A+randomized,+controlled+trial&author=Kim,+J.Y.&author=Gum,+S.N.&author=Paik,+J.K.&author=Lim,+H.H.&author=Kim,+K.C.&author=Ogasawara,+K.&author=Inoue,+K.&author=Park,+S.&author=Jang,+Y.&author=Lee,+J.H.&publication\_year=2008&journal=Hypertens.+Res.&volume=31&pages=1583%E2%80%931588&doi=10.1291/hypres.31.1583]** [CrossRef (<https://doi.org/10.1291/hypres.31.1583>)]

12. Wu, C.; Gao, C.; Lü, S.; Xu, X.; Wen, N.; Zhang, S.; Liu, M. Construction of polylysine dendrimer nanocomposites carrying nattokinase and their application in thrombolysis. *J. Biomed. Mater. Res. A* **2018**, *106*, 440–449. [Google Scholar (https://scholar.google.com/scholar\_lookup?title=Construction+of+polylysine+dendrimer+nanocomposites+carrying+nattokinase+and+their+application+in+thrombolysis&author=Wu,+C.&author=Gao,+C.&author=L%C3%BC,+S.&author=Xu,+X.&author=Wen,+N.&author=Zhang,+S.&author=Liu,+M.&publication\_year=2018&journal=J.+Biomed.+Mater.+Res.+A&volume=106&pages=440%E2%80%93449&doi=10.1002/jbm.a.36232)] [CrossRef (https://doi.org/10.1002/jbm.a.36232)]
13. Lampe, B.J.; English, J.C. Toxicological assessment of nattokinase derived from *Bacillus subtilis* var *natto*. *Food Chem. Toxicol.* **2016**, *88*, 87–99. [Google Scholar (https://scholar.google.com/scholar\_lookup?title=Toxicological+assessment+of+nattokinase+derived+from+Bacillus+subtilis+var+natto&author=Lampe,+B.J.&author=English,+J.C.&publication\_year=2016&journal=Food+Chem.+Toxicol.&volume=88&pages=87%E2%80%9399&doi=10.1016/j.fct.2015.12.025)] [CrossRef (https://doi.org/10.1016/j.fct.2015.12.025)]
14. Kou, Y.; Feng, R.; Chen, J.; Duan, L.; Wang, S.; Hu, Y.; Zhang, N.; Wang, T.; Deng, Y.; Song, Y. Development of a nattokinase-polysialic acid complex for advanced tumor treatment. *Eur. J. Pharm. Sci.* **2020**, *145*, 105241. [Google Scholar (https://scholar.google.com/scholar\_lookup?title=Development+of+a+nattokinase-polysialic+acid+complex+for+advanced+tumor+treatment&author=Kou,+Y.&author=Feng,+R.&author=Chen,+J.&author=Duan,+L.&author=Wang,+S.&author=Hu,+Y.&author=Zhang,+N.&author=Wang,+T.&author=Deng,+Y.&author=Song,+Y.&publication\_year=2020&journal=Eur.+J.+Pharm.+Sci.&volume=145&pages=105241&doi=10.1016/j.ejps.2020.105241)] [CrossRef (https://doi.org/10.1016/j.ejps.2020.105241)]
15. Yan, Y.; Wang, Y.; Qian, J.; Wu, S.; Ji, Y.; Liu, Y.; Zeng, J.; Gong, A. Nattokinase Crude Extract Inhibits Hepatocellular Carcinoma Growth in Mice. *J. Microbiol. Biotechnol.* **2019**, *29*, 1281–1287. [Google Scholar (https://scholar.google.com/scholar\_lookup?title=Nattokinase+Crude+Extract+Inhibits+Hepatocellular+Carcinoma+Growth+in+Mice&author=Yan,+Y.&author=Wang,+Y.&author=Qian,+J.&author=Wu,+S.&author=Ji,+Y.&author=Liu,+Y.&author=Zeng,+J.&author=Gong,+A.&publication\_year=2019&journal=J.+Microbiol.+Biotechnol.&volume=29&pages=1281%E2%80%931287&doi=10.4014/jmb.1812.12058)] [CrossRef (https://doi.org/10.4014/jmb.1812.12058)]

16. Oba, M.; Rongduo, W.; Saito, A.; Okabayashi, T.; Yokota, T.; Yasuoka, J.; Sato, Y.; Nishifuji, K.; Wake, H.; Nibu, Y.; et al. Natto extract, a Japanese fermented soybean food, directly inhibits viral infections including SARS-CoV-2 in vitro. *Biochem. Biophys. Res. Commun.* **2021**, *570*, 21–25. [Google Scholar (https://scholar.google.com/scholar\_lookup?title=Natto+extract,+a+Japanese+fermented+soybean+food,+directly+inhibits+viral+infections+including+SARS-CoV-2+in+vitro&author=Oba,+M.&author=Rongduo,+W.&author=Saito,+A.&author=Okabayashi,+T.&author=Yokota,+T.&author=Yasuoka,+J.&author=Sato,+Y.&author=Nishifuji,+K.&author=Wake,+H.&author=Nibu,+Y.&publication\_year=2021&journal=Biochem.+Biophys.+Res.+Commun.&volume=570&pages=21%E2%80%9325&doi=10.1016/j.bbrc.2021.07.034)] [CrossRef (https://doi.org/10.1016/j.bbrc.2021.07.034)]
17. Smaoui, M.R.; Yahyaoui, H. Unraveling the stability landscape of mutations in the SARS-CoV-2 receptor-binding domain. *Sci. Rep.* **2021**, *11*, 9166. [Google Scholar (https://scholar.google.com/scholar\_lookup?title=Unraveling+the+stability+landscape+of+mutations+in+the+SARS-CoV-2+receptor-binding+domain&author=Smaoui,+M.R.&author=Yahyaoui,+H.&publication\_year=2021&journal=Sci.+Rep.&volume=11&pages=9166&doi=10.1038/s41598-021-88696-5)] [CrossRef (https://doi.org/10.1038/s41598-021-88696-5)]
18. Carter, P.; Wells, J.A. Dissecting the catalytic triad of a serine protease. *Nature* **1988**, *332*, 564–568. [Google Scholar (https://scholar.google.com/scholar\_lookup?title=Dissecting+the+catalytic+triad+of+a+serine+protease&author=Carter,+P.&author=Wells,+J.A.&publication\_year=1988&journal=Nature&volume=332&pages=564%E2%80%93568&doi=10.1038/332564a0)] [CrossRef (https://doi.org/10.1038/332564a0)]
19. Yanagisawa, Y.; Chatake, T.; Naito, S.; Ohsugi, T.; Yatagai, C.; Sumi, H.; Kawaguchi, A.; Chiba-Kamosida, K.; Ogawa, M.; Adachi, T.; et al. X-ray structure determination and deuteration of nattokinase. *J. Synchrotron Radiat.* **2013**, *20 Pt 6*, 875–879. [Google Scholar (https://scholar.google.com/scholar\_lookup?title=X-ray+structure+determination+and+deuteration+of+nattokinase&author=Yanagisawa,+Y.&author=Chatake,+T.&author=Naito,+S.&author=Ohsugi,+T.&author=Yatagai,+C.&author=Sumi,+H.&author=Kawaguchi,+A.&author=Chiba-Kamosida,+K.&author=Ogawa,+M.&author=Adachi,+T.&publication\_year=2013&journal=J.+Synchrotron+Radiat.&volume=20&issue=Pt+6&pages=875%E2%80%93879&doi=10.1107/S0909049513020700)] [CrossRef (https://doi.org/10.1107/S0909049513020700)]



Chen, H.; McGowan, E.M.; Ren, N.; Lal, S.; Nassif, N.; Shad-Kaneez, F.; Qu, X.; Lin, Y. Nattokinase: A Promising Alternative in Prevention and Treatment of Cardiovascular Diseases. *Biomark. Insights* **2018**, *13*, 1177271918785130. [Google Scholar ([https://scholar.google.com/scholar\\_lookup?title=Nattokinase:+A+Promising+Alternative+in+Prevention+and+Treatment+of+Cardiovascular+Diseases&author=Chen,+H.&author=McGowan,+E.M.&author=Ren,+N.&author=Lal,+S.&author=Nassif,+N.&author=Shad-Kaneez,+F.&author=Qu,+X.&author=Lin,+Y.&publication\\_year=2018&journal=Biomark.+Insights&volume=13&pages=1177271918785130&doi=10.1177/1177271918785130](https://scholar.google.com/scholar_lookup?title=Nattokinase:+A+Promising+Alternative+in+Prevention+and+Treatment+of+Cardiovascular+Diseases&author=Chen,+H.&author=McGowan,+E.M.&author=Ren,+N.&author=Lal,+S.&author=Nassif,+N.&author=Shad-Kaneez,+F.&author=Qu,+X.&author=Lin,+Y.&publication_year=2018&journal=Biomark.+Insights&volume=13&pages=1177271918785130&doi=10.1177/1177271918785130))] [CrossRef (<https://doi.org/10.1177/1177271918785130>)] [Green Version (<https://journals.sagepub.com/doi/pdf/10.1177/1177271918785130>)]

21. Cancarevic, I.; Malik, B.H. SARS-CoV-2 (COVID 19) Infection in Hypertensive Patients and in Patients with Cardiac Disease. *Cureus* **2020**, *12*, e8557. [Google Scholar ([https://scholar.google.com/scholar\\_lookup?title=SARS-CoV-2+\(COVID+19\)+Infection+in+Hypertensive+Patients+and+in+Patients+with+Cardiac+Disease&author=Cancarevic,+I.&author=Malik,+B.H.&publication\\_year=2020&journal=Cureus&volume=12&pages=e8557&doi=10.7759/cureus.8557](https://scholar.google.com/scholar_lookup?title=SARS-CoV-2+(COVID+19)+Infection+in+Hypertensive+Patients+and+in+Patients+with+Cardiac+Disease&author=Cancarevic,+I.&author=Malik,+B.H.&publication_year=2020&journal=Cureus&volume=12&pages=e8557&doi=10.7759/cureus.8557))] [CrossRef (<https://doi.org/10.7759/cureus.8557>)]

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

© 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/> (<https://creativecommons.org/licenses/by/4.0/>)).

## Share and Cite



(mailto:?)

&subject=From%20MDPI%3A%20%22Degradative%20Effect%20of%20Nattokinase%20on%20Spike%20Protein%20of%20SARS-CoV-2" &body=https://www.mdpi.com/1792976%3A%0A%0ADegradative%20Effect%20of%20Nattokinase%20on%20Spike%20Protein%20of%20SARS-CoV-2%0A%0AAbstract%3A%20The%20coronavirus%20disease%202019%20%28COVID-19%29%2C%20caused%20by%20the%20severe%20acute%20respiratory%20syndrome%20coronavirus%202%20%28SARS-CoV-2%29%2C%20emerged%20as%20a%20pandemic%20and%20has%20inflicted%20enormous%20damage%20on%20the%20lives%20of%20the%20people%20and%20economy

MDPI (i) of many countries worldwide. However, therapeutic agents against SARS-CoV-2 remain unclear. SARS-CoV-

2 has a spike protein and cleavage of the S protein is essential for viral entry. Nattokinase is produced by *Bacillus subtilis* var. *natto* and is beneficial to human health. In this study, we examined the effect of nattokinase on the S protein of SARS-CoV-

2. When cell lysates transfected with S protein were incubated with nattokinase, the S protein was degraded in a dose- and time-

dependent manner. Immunofluorescence analysis showed that S protein on the cell surface was degraded when nattokinase was added to the culture medium. Thus, our findings suggest that nattokinase exhibits potential for the inhibition of SARS-CoV-

2 infection via S protein degradation.) 

([https://twitter.com/intent/tweet?](https://twitter.com/intent/tweet?text=Degradative+Effect+of+Nattokinase+on+Spike+Protein+of+SARS-CoV-2&hashtags=mdpimolecules&url=https%3A%2F%2Fwww.mdpi.com%2F1792976&via=Molecules_MDPI)

[text=Degradative+Effect+of+Nattokinase+on+Spike+Protein+of+SARS-CoV-](https://twitter.com/intent/tweet?text=Degradative+Effect+of+Nattokinase+on+Spike+Protein+of+SARS-CoV-2&hashtags=mdpimolecules&url=https%3A%2F%2Fwww.mdpi.com%2F1792976&via=Molecules_MDPI)

[2&hashtags=mdpimolecules&url=https%3A%2F%2Fwww.mdpi.com%2F1792976&via=](https://twitter.com/intent/tweet?text=Degradative+Effect+of+Nattokinase+on+Spike+Protein+of+SARS-CoV-2&hashtags=mdpimolecules&url=https%3A%2F%2Fwww.mdpi.com%2F1792976&via=Molecules_MDPI)

[Molecules\\_MDPI](https://twitter.com/intent/tweet?text=Degradative+Effect+of+Nattokinase+on+Spike+Protein+of+SARS-CoV-2&hashtags=mdpimolecules&url=https%3A%2F%2Fwww.mdpi.com%2F1792976&via=Molecules_MDPI)  ([http://www.linkedin.com/shareArticle?](http://www.linkedin.com/shareArticle?mini=true&url=https%3A%2F%2Fwww.mdpi.com%2F1792976&title=Degradative%20Effect%20of%20Nattokinase%20on%20Spike%20Protein%20of%20SARS-CoV-2%26source%3Dhttps%3A%2F%2Fwww.mdpi.com%26summary%3DThe%20coronavirus%20disease%202019%20%28COVID-19%29%2C%20caused%20by%20the%20severe%20acute%20respiratory%20syndrome%20coronavirus%202020%28SARS-CoV-2%29%2C%20emerged%20as%20a%20pandemic%20and%20has%20inflicted%20enormous%20damage%20on%20the%20lives%20of%20the%20people%20and%20economy%20of%20many%20countries%20worldwide.%20However%2C%20%5B...%5D)

[mini=true&url=https%3A%2F%2Fwww.mdpi.com%2F1792976&title=Degradative%20Eff](https://twitter.com/intent/tweet?text=Degradative+Effect+of+Nattokinase+on+Spike+Protein+of+SARS-CoV-2&hashtags=mdpimolecules&url=https%3A%2F%2Fwww.mdpi.com%2F1792976&via=Molecules_MDPI)

[ect%20of%20Nattokinase%20on%20Spike%20Protein%20of%20SARS-CoV-](https://twitter.com/intent/tweet?text=Degradative+Effect+of+Nattokinase+on+Spike+Protein+of+SARS-CoV-2&hashtags=mdpimolecules&url=https%3A%2F%2Fwww.mdpi.com%2F1792976&via=Molecules_MDPI)

[2%26source%3Dhttps%3A%2F%2Fwww.mdpi.com%26summary%3DThe%20coronavir](https://twitter.com/intent/tweet?text=Degradative+Effect+of+Nattokinase+on+Spike+Protein+of+SARS-CoV-2&hashtags=mdpimolecules&url=https%3A%2F%2Fwww.mdpi.com%2F1792976&via=Molecules_MDPI)

[us%20disease%202019%20%28COVID-](https://twitter.com/intent/tweet?text=Degradative+Effect+of+Nattokinase+on+Spike+Protein+of+SARS-CoV-2&hashtags=mdpimolecules&url=https%3A%2F%2Fwww.mdpi.com%2F1792976&via=Molecules_MDPI)

[19%29%2C%20caused%20by%20the%20severe%20acute%20respiratory%20syndrome](https://twitter.com/intent/tweet?text=Degradative+Effect+of+Nattokinase+on+Spike+Protein+of+SARS-CoV-2&hashtags=mdpimolecules&url=https%3A%2F%2Fwww.mdpi.com%2F1792976&via=Molecules_MDPI)

[%20coronavirus%202020%28SARS-CoV-](https://twitter.com/intent/tweet?text=Degradative+Effect+of+Nattokinase+on+Spike+Protein+of+SARS-CoV-2&hashtags=mdpimolecules&url=https%3A%2F%2Fwww.mdpi.com%2F1792976&via=Molecules_MDPI)

[2%29%2C%20emerged%20as%20a%20pandemic%20and%20has%20inflicted%20enor](https://twitter.com/intent/tweet?text=Degradative+Effect+of+Nattokinase+on+Spike+Protein+of+SARS-CoV-2&hashtags=mdpimolecules&url=https%3A%2F%2Fwww.mdpi.com%2F1792976&via=Molecules_MDPI)

[mous%20damage%20on%20the%20lives%20of%20the%20people%20and%20economy](https://twitter.com/intent/tweet?text=Degradative+Effect+of+Nattokinase+on+Spike+Protein+of+SARS-CoV-2&hashtags=mdpimolecules&url=https%3A%2F%2Fwww.mdpi.com%2F1792976&via=Molecules_MDPI)

[%20of%20many%20countries%20worldwide.%20However%2C%20%5B...%5D](https://twitter.com/intent/tweet?text=Degradative+Effect+of+Nattokinase+on+Spike+Protein+of+SARS-CoV-2&hashtags=mdpimolecules&url=https%3A%2F%2Fwww.mdpi.com%2F1792976&via=Molecules_MDPI) 

(<https://www.facebook.com/sharer.php?u=https://www.mdpi.com/1792976>) 

(<http://www.reddit.com/submit?url=https://www.mdpi.com/1792976>)  
(<http://www.mendeley.com/import/?url=https://www.mdpi.com/1792976>)

## MDPI and ACS Style

Tanikawa, T.; Kiba, Y.; Yu, J.; Hsu, K.; Chen, S.; Ishii, A.; Yokogawa, T.; Suzuki, R.; Inoue, Y.; Kitamura, M. Degradative Effect of Nattokinase on Spike Protein of SARS-CoV-2. *Molecules* **2022**, *27*, 5405. <https://doi.org/10.3390/molecules27175405>

### Chicago/Turabian Style

Tanikawa, Takashi, Yuka Kiba, James Yu, Kate Hsu, Shinder Chen, Ayako Ishii, Takami Yokogawa, Ryuichiro Suzuki, Yutaka Inoue, and Masashi Kitamura. 2022. "Degradative Effect of Nattokinase on Spike Protein of SARS-CoV-2" *Molecules* 27, no. 17: 5405. <https://doi.org/10.3390/molecules27175405>

### Find Other Styles

▼ type a publisher, journal or format name


### Article Metrics

#### Citazioni

Rif. incrociato	Scopo	Web della scienza	PMC	PubMed	Google Scholar
2	2 ( <a href="https://partnerID=HzO">https://partnerID=HzO</a> )	GWVersion=28	( <a href="https://www.ndinkname=pubr">https://www.ndinkname=pubr</a> )	1 ( <a href="http">http</a> )	2&volume=27&

### Article Access Statistics

Article access statistics



● Article Views

For more information on the journal statistics, click [here \(/journal/molecules/stats\)](/journal/molecules/stats).



Multiple requests from the same IP address are counted as one view.

 [\(/toggle desktop layout cookie\)](#)  

***Molecules*** (</journal/molecules>), EISSN 1420-3049, Published by MDPI

[RSS \(/rss/journal/molecules\)](/rss/journal/molecules)    [Content Alert \(/journal/molecules/toc-alert\)](/journal/molecules/toc-alert)

#### Further Information

[Article Processing Charges \(/apc\)](/apc)

[Pay an Invoice \(/about/payment\)](/about/payment)

[Open Access Policy \(/openaccess\)](/openaccess)

[Contact MDPI \(/about/contact\)](/about/contact)

[Jobs at MDPI \(https://careers.mdpi.com\)](https://careers.mdpi.com)

#### Guidelines

[For Authors \(/authors\)](/authors)

[For Reviewers \(/reviewers\)](/reviewers)

[For Editors \(/editors\)](/editors)

[For Librarians \(/librarians\)](/librarians)

[For Publishers \(/publishing\\_services\)](/publishing_services)

[For Societies \(/societies\)](/societies)

[For Conference Organizers \(/conference\\_organizers\)](/conference_organizers)

MDPI Initiatives

[Sciforum \(https://sciforum.net\)](https://sciforum.net)

[MDPI Books \(https://www.mdpi.com/books\)](https://www.mdpi.com/books)

[Preprints.org \(https://www.preprints.org\)](https://www.preprints.org)

[Scilit \(https://www.scilit.net\)](https://www.scilit.net)

[SciProfiles \(https://sciprofiles.com\)](https://sciprofiles.com)

[Encyclopedia \(https://encyclopedia.pub\)](https://encyclopedia.pub)

[JAMS \(https://jams.pub\)](https://jams.pub)

[Proceedings Series \(/about/proceedings\)](/about/proceedings)


#### Follow MDPI

[LinkedIn \(https://www.linkedin.com/company/mdpi\)](https://www.linkedin.com/company/mdpi)

[Facebook \(https://www.facebook.com/MDPIOpenAccessPublishing\)](https://www.facebook.com/MDPIOpenAccessPublishing)



**Iscriviti per ricevere le notifiche di  
rilascio dei numeri e le newsletter  
dalle riviste MDPI**

Seleziona le opzioni 

Enter your email address...

**sottoscrivi**

© 1996-2023 MDPI (Basilea, Svizzera) se non diversamente specificato

**[Disclaimer](#) [Termini e Condizioni \(/about/terms-and-conditions\)](#),  
[politica sulla riservatezza \(/about/privacy\)](#)**